14.2 Sepsis and Septic Shock*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I feel very unwell and shivery." **Objective** • Temperature 39.3 °C (or below 36 °C) • BP 85/45 mmHg, HR 128/min, RR 28/min • Lactate 4 mmol/L • New confusion	**ACTUAL** Ineffective peripheral tissue perfusion related to vasodilation and capillary leak as evidenced by hypotension, raised lactate, mottled skin and confusion.	• MAP of 65 mmHg or more and lactate falling within 6 hours.	• Follow the sepsis bundle: blood cultures first, then IV antibiotics within 1 hour. • Give an IV fluid bolus as prescribed and reassess. • Measure lactate and repeat as ordered. • Start vasopressors as prescribed if hypotension persists. • Measure urine output hourly.	• Goal reached when MAP is on target and lactate is coming down.
Subjective *No specific complaint* **Objective** • Temperature 39.3 °C • Rigors	**ACTUAL** Hyperthermia related to infection as evidenced by a temperature of 39.3 °C and rigors.	• Temperature below 38 °C within 6 hours.	• Give the prescribed antipyretic. • Remove extra covers. • Measure temperature hourly.	• Met as temperature falls.
Subjective *No specific complaint* **Objective** • Septic shock • Hypotension	**RISK** Risk for decreased cardiac output as evidenced by risk factors: septic cardiomyopathy and dysrhythmias.	• Rhythm and cardiac output stable.	• Keep him on continuous ECG monitoring. • Follow electrolytes. • Report new dysrhythmias.	• Achieved while he remains stable.

*References for this care plan: [20], [35], [128]. Full entries are listed in the References section of the CPMN book.