15.17 Normal Postpartum Care*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "My stitches hurt when I sit." • Rates pain 5/10 **Objective** • Repaired perineal tear • Slight swelling	**ACTUAL** Acute pain related to perineal trauma as evidenced by pain on sitting and swelling.	• Pain at or below 3/10.	• Apply a cold pack to the perineum in the first 24 hours. • Give analgesia as prescribed. • Teach perineal hygiene and frequent pad changes.	• Considered met when pain is controlled.
Subjective • "How do I care for the baby's cord and bath?" **Objective** • First baby • Unsure about bathing	**ACTUAL** Inadequate health knowledge related to newborn care and self-care as evidenced by questioning.	• Shows basic newborn care before discharge.	• Teach cord care, bathing, feeding cues and safe sleep. • Teach the baby's danger signs: poor feeding, jaundice, fever, difficulty breathing. • Teach her own warning signs: heavy bleeding, fever, leg pain, low mood.	• Goal reached when she demonstrates it.
Subjective *No specific complaint* **Objective** • Recent birth	**RISK** Risk for excessive bleeding as evidenced by risk factors: postpartum uterus.	• Fundus firm; lochia normal.	• Check fundus and lochia regularly in the first 24 hours. • Encourage her to empty her bladder, since a full bladder stops the uterus contracting well. • Teach fundal massage.	• Met if no hemorrhage occurs.

* **References for this care plan:** [35], [47], [96], [143]. Full entries are listed in the References section of the CPMN book.