16.8 Neonatal Hypothermia*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective *No specific complaint* **Objective** • Temperature 35.5 °C • Cool hands and feet • Poor feeding	**ACTUAL** Decreased body temperature related to heat loss and immature thermoregulation as evidenced by a temperature of 35.5 °C.	• Temperature 36.5–37.5 °C within 2 hours.	• Rewarm skin-to-skin or in an incubator. • Remove wet clothing. • Check temperature every 30 minutes until normal. • Check blood glucose.	• Outcome reached when temperature is normal.
Subjective *No specific complaint* **Objective** • Lethargic • Weak suck	**ACTUAL** Ineffective infant feeding dynamics related to cold stress as evidenced by poor feeding.	• Feeds well again once warm.	• Feed once temperature is normal. • Support breastfeeding or give expressed milk. • Record intake.	• Considered met when feeding resumes.
Subjective *No specific complaint* **Objective** • Cold stress • Lethargic	**RISK** Risk for physical injury as evidenced by risk factors: hypoglycemia associated with increased glucose use during cold stress.	• Blood glucose stays normal.	• Check glucose. • Feed early. • Give IV dextrose if ordered.	• Goal reached while glucose is normal.

*References for this care plan: [35], [38], [142]. Full entries are listed in the References section of the CPMN book.