17.16 Acute Lymphoblastic Leukemia in Children*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • Child says, "I feel sick after the medicine." **Objective** • Vomits after chemotherapy • Eating less	**ACTUAL** Ineffective nausea self-management related to chemotherapy as evidenced by vomiting and reduced intake.	• Less nausea and enough intake.	• Give antiemetics before chemotherapy. • Offer foods she likes in small amounts. • Follow her weight.	• Considered met as nausea lessens.
Subjective • Child says, "I do not want another needle." **Objective** • Cries during procedures • Clings to her mother	**ACTUAL** Excessive fear related to painful procedures as evidenced by crying and resistance.	• Copes with procedures.	• Apply a topical anesthetic before needles. • Prepare her through therapeutic play. • Let a parent stay.	• Goal reached when she is coping.
Subjective *No specific complaint* **Objective** • Neutropenia	**RISK** Risk for infection as evidenced by risk factors: chemotherapy-induced neutropenia.	• No infection; any fever treated within 1 hour.	• Put protective measures in place. • Teach the parents to report a temperature of 38 °C or more. • Avoid live vaccines, as advised.	• Met if no infection develops.

*References for this care plan: [22], [35], [38], [63]. Full entriFull entries are listed in the References section of the CPMN book.