## 2.18 Nausea and Vomiting*

| Assessment | Nursing Diagnosis | Plan / Expected Outcome | Nursing Interventions | Evaluation |
|---|---|---|---|---|
| **Subjective**<br>• "I feel sick all the time."<br>**Objective**<br>• Vomited four times in 6 hours<br>• Pale, sweaty | **ACTUAL**<br>Ineffective nausea self-management related to gastric irritation and pharmaceutical agents as evidenced by reported nausea and vomiting. | • Nausea eases within 2 hours. | • Give the prescribed antiemetic.<br>• Remove odors and air the room.<br>• Offer small amounts of clear fluid once tolerated.<br>• Provide mouth care after vomiting. | • Considered met when nausea lessens and vomiting stops. |
| **Subjective**<br>• Thirsty<br>**Objective**<br>• Dry mouth<br>• Passing less urine<br>• HR 108/min | **ACTUAL**<br>Inadequate fluid volume related to vomiting as evidenced by dry mucosa and reduced urine output. | • Adequate urine output within 8 hours. | • Chart intake, output and urine volume every 4 hours.<br>• Give IV fluids as prescribed.<br>• Observe vital signs and capillary refill.<br>• Introduce sips of clear fluid as the vomiting settles. | • Goal reached with adequate output. |
| **Subjective**<br>*No specific complaint*<br>**Objective**<br>• Repeated vomiting | **RISK**<br>Risk for impaired water-electrolyte balance as evidenced by risk factors: vomiting. | • Electrolytes within normal limits. | • Follow serum electrolytes.<br>• Report abnormal values.<br>• Replace as ordered. | • Met if values remain normal. |

* **References for this care plan:** [35], [37], [48], [64], [130]. Full entries are listed in the References section of the CPMN book.