3.17 Obstructive Sleep Apnea*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • Wife reports loud snoring and pauses in breathing • "I feel sleepy all day." **Objective** • BMI 36 kg/m² • Large neck circumference • Oxygen desaturation at night	**ACTUAL** Ineffective sleep pattern related to upper airway obstruction during sleep as evidenced by snoring, apneas and daytime sleepiness.	• Uses CPAP and reports better sleep within 2 weeks.	• Teach mask fitting and care of the CPAP equipment. • Encourage weight loss. • Advise no alcohol or sedatives before bed. • Suggest sleeping on his side.	• Considered met when CPAP is used every night.
Subjective • "I fell asleep while driving." **Objective** • Drowsy by day • Dozes off during the interview	**ACTUAL** Excessive fatigue burden related to poor sleep quality as evidenced by daytime sleepiness.	• Feels more alert by day within 2 weeks of starting CPAP.	• Advise him not to drive or use machinery while drowsy, until treated. • Encourage regular sleep times. • Repeat the advice on evening sedatives and alcohol. • Reassess alertness at follow-up.	• Goal reached when he reports better alertness.
Subjective *No specific complaint* **Objective** • Repeated nocturnal hypoxemia • Obesity	**RISK** Risk for impaired cardiovascular function as evidenced by risk factors: nocturnal hypoxemia and hypertension.	• BP on target with nightly CPAP.	• Check BP at every visit. • Encourage nightly CPAP and regular antihypertensive use. • Encourage weight loss and regular exercise. • Teach him to report chest pain or palpitations.	• Met when BP is controlled.

*References for this care plan: [32], [35], [37], [132]. Full entries are listed in the References section of the CPMN book.