5.14 Myasthenia Gravis*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "My eyelids droop and I get tired chewing." **Objective** • Ptosis • Weak chewing and swallowing	**ACTUAL** Impaired swallowing related to neuromuscular weakness as evidenced by difficulty chewing and swallowing.	• Eats safely without choking.	• Give the anticholinesterase 30–45 minutes before meals, as prescribed. • Offer soft foods and let her rest during meals. • Keep suction ready.	• Outcome reached when eating is safe.
Subjective *No specific complaint* **Objective** • Tiredness builds through the day • Ptosis worse in the evening	**ACTUAL** Excessive fatigue burden related to muscle weakness as evidenced by weakness that worsens with activity.	• Plans activities for the times when she is strongest.	• Plan activities soon after a dose. • Encourage rest periods. • Teach ways to save energy.	• Considered met when activities are planned this way.
Subjective *No specific complaint* **Objective** • Weak respiratory muscles • Weak cough	**RISK** Risk for aspiration as evidenced by risk factors: impaired swallowing and weak cough.	• No aspiration; any crisis is recognized early.	• Watch breathing and swallowing. • Teach the signs of myasthenic and cholinergic crisis.	• Goal reached if aspiration does not occur.

* **References for this care plan:** [32], [35], [37]. Full entries are listed in the References section of the CPMN book.