7.14 Urinary Retention*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I need to pass urine but nothing comes." **Objective** • Distended bladder, dull to percussion • Bladder scan 700 mL • On an anticholinergic drug	**ACTUAL** Impaired urinary elimination related to bladder outlet obstruction or medication effect as evidenced by bladder distension and inability to void.	• Bladder emptied and voiding on his own within 24 hours.	• Help him void: privacy, running water, standing or sitting. • Catheterize as ordered and record the volume. • Review anticholinergic and opioid drugs with the physician. • Do a trial without catheter as ordered and scan for residual.	• Goal reached when he voids spontaneously with a small residual.
Subjective • Rates lower abdominal discomfort 7/10 **Objective** • Restless • Sweating	**ACTUAL** Acute pain related to bladder distension as evidenced by discomfort and restlessness.	• Relief after drainage.	• Drain the bladder as ordered. • Reassess pain afterward. • Watch for hematuria after a large drainage.	• Met once he is relieved.
Subjective *No specific complaint* **Objective** • Catheterized • Residual urine	**RISK** Risk for infection as evidenced by risk factors: urinary catheter and urinary stasis.	• No UTI.	• Insert the catheter aseptically. • Keep the drainage system closed. • Remove the catheter as early as possible.	• Achieved if no infection develops.

*References for this care plan: [28], [32], [35], [37]. Full entries are listed in the References section of the CPMN book.