## 7.2 Pyelonephritis*

| Assessment | Nursing Diagnosis | Plan / Expected Outcome | Nursing Interventions | Evaluation |
|---|---|---|---|---|
| **Subjective**<br>• "I have pain in my side and I feel shivery."<br>**Objective**<br>• Temperature 39 °C<br>• Costovertebral angle tenderness<br>• Pus cells in urine | **ACTUAL**<br>Hyperthermia related to renal infection as evidenced by a temperature of 39 °C and chills. | • Afebrile within 72 hours of antibiotics. | • Send urine and blood cultures as ordered.<br>• Give IV antibiotics on time.<br>• Give antipyretics.<br>• Encourage drinking, or give IV fluids as ordered. | • Considered met once afebrile. |
| **Subjective**<br>• Rates flank pain 7/10<br>**Objective**<br>• Guarding<br>• Restless | **ACTUAL**<br>Acute pain related to renal inflammation as evidenced by flank pain. | • Pain at or below 3/10 within 24 hours. | • Give analgesia as prescribed.<br>• Apply a warm compress to the flank.<br>• Encourage fluids.<br>• Reassess pain regularly. | • Goal reached when pain is controlled. |
| **Subjective**<br>*No specific complaint*<br>**Objective**<br>• Renal infection<br>• Tachycardia | **RISK**<br>Risk for shock as evidenced by risk factors: possible urosepsis. | • Vital signs and urine output stay stable. | • Take vital signs every 4 hours and follow urine output.<br>• Report hypotension, tachycardia or confusion.<br>• Give IV antibiotics on time. | • Met while she remains stable. |

* **References for this care plan:** [20], [32], [35], [37]. Full entries are listed in the References section of the CPMN book.