10.9 Hemophilia*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "My knee swelled after a small fall." • Rates pain 7/10 **Objective** • Bleeding into the knee joint • Factor VIII deficiency • Knee movement limited	**ACTUAL** Acute pain related to intra-articular bleeding as evidenced by swelling, pain and limited movement.	• Bleeding stopped and pain at or below 3/10 within 24 hours.	• Give factor concentrate immediately, as prescribed. • Apply rest, ice, compression and elevation. • Give paracetamol or an opioid as prescribed; no aspirin or NSAIDs. • Avoid IM injections.	• Goal reached when pain is controlled and swelling goes down.
Subjective • "How do I give the factor at home?" **Objective** • Home therapy planned • Cannot describe reconstitution	**ACTUAL** Inadequate health knowledge related to home factor replacement as evidenced by questioning.	• Patient or caregiver prepares and gives the factor IV correctly.	• Teach reconstitution and IV administration. • Teach recording of doses and bleeds. • Teach when to seek urgent help, such as after a head injury or major bleed.	• Met when demonstrated correctly.
Subjective *No specific complaint* **Objective** • Severe factor deficiency	**RISK** Risk for excessive bleeding as evidenced by risk factors: clotting factor deficiency.	• No major bleeding episodes.	• Suggest safe exercise such as swimming; avoid contact sports. • Advise wearing medical identification. • Teach oral hygiene to prevent gum bleeding.	• Achieved if no major bleed occurs.

*References for this care plan: [32], [35], [38]. Full entries are listed in the References section of the CPMN book.