14.6 Multiple Trauma*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I feel weak and dizzy." **Objective** • Pelvic and femoral fractures • BP 86/54 mmHg, HR 128/min • Pale, cold skin	**ACTUAL** Inadequate fluid volume related to hemorrhage from multiple injuries as evidenced by hypotension, tachycardia and cold skin.	• MAP of 65 mmHg or more; urine at least 0.5 mL/kg/hour.	• Carry out the primary survey: airway with cervical spine control, breathing, circulation. • Stop external bleeding and apply a pelvic binder as ordered. • Give warmed blood products per the massive transfusion protocol. • Take vital signs every 5–15 minutes and measure urine hourly. • Prepare for imaging and surgery.	• Achieved when hemodynamics are stable.
Subjective • "Everything hurts." • Rates pain 9/10 **Objective** • Multiple fractures • Grimacing; guarding	**ACTUAL** Acute pain related to multiple tissue injuries as evidenced by a pain score of 9/10 and guarding.	• Pain bearable within 1 hour.	• Give IV analgesia in small, titrated doses as prescribed. • Splint the fractures. • Check pain and sedation often. • Handle and move him gently.	• Outcome reached when pain is tolerable.
Subjective *No specific complaint* **Objective** • Open wounds • Invasive lines and drains	**RISK** Risk for infection as evidenced by risk factors: open wounds, contamination and invasive devices.	• No wound infection.	• Clean and dress wounds aseptically. • Give tetanus prophylaxis and antibiotics as ordered. • Watch temperature and the wounds.	• Considered met if no infection develops.

* **References for this care plan:** [2], [35], [128]. Full entries are listed in the References section of the CPMN book.