15.11 Anemia in Pregnancy*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I feel tired and dizzy." **Objective** • Hb 9 g/dL • Pale	**ACTUAL** Excessive fatigue burden related to anemia as evidenced by tiredness and dizziness.	• More energy within 4 weeks of treatment.	• Give iron as prescribed. • Plan rest periods. • Follow hemoglobin. • Teach iron-rich foods.	• Met as energy improves.
Subjective • "I stopped iron because of constipation." **Objective** • Not taking iron • Hb not improving	**ACTUAL** Ineffective health self-management related to adverse effects of iron as evidenced by discontinuation of treatment.	• Takes iron again as prescribed.	• Manage side effects with fluid and fiber. • Explain why iron matters for her and the baby. • Discuss a different preparation with the physician.	• Achieved once iron is restarted.
Subjective *No specific complaint* **Objective** • Hb 9 g/dL at 36 weeks	**RISK** Risk for excessive bleeding as evidenced by risk factors: reduced reserve to tolerate blood loss at birth.	• Hemoglobin better before delivery.	• Follow hemoglobin. • Plan care in labor with the obstetric team. • Make sure blood is available at delivery if needed.	• Outcome reached as Hb improves.

*References for this care plan: [35], [47], [95], [138]. Full entries are listed in the References section of the CPMN book.