15.8 Eclampsia*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective *No specific complaint* **Objective** • Generalized seizure • BP 170/110 mmHg • Frothy secretions at the mouth	**ACTUAL** Ineffective airway clearance related to seizure activity as evidenced by frothy secretions.	• Airway clear and seizure controlled.	• Call for help and turn her onto her left side. • Keep the airway open; suction and give oxygen. • Give magnesium sulfate as prescribed. • Do not restrain her; protect her from injury.	• Achieved when the seizure is controlled.
Subjective • Husband says, “She does not know where she is.” **Objective** • Drowsy after the seizure	**ACTUAL** Acute confusion related to the postictal state as evidenced by drowsiness and disorientation.	• Oriented within hours of the seizure.	• Do neurological checks every 15–30 minutes. • Reorient her gently. • Observe her closely with the rails raised.	• Outcome reached once she is oriented.
Subjective *No specific complaint* **Objective** • Maternal seizure	**RISK** Risk for impaired maternal-fetal dyad as evidenced by risk factors: maternal hypoxia during seizures.	• Fetal well-being maintained until a safe delivery.	• Keep the fetal heart on continuous monitoring. • Give the mother oxygen. • Prepare for delivery once she is stable.	• Considered met when the outcome is safe.

* **References for this care plan:** [1], [35], [47], [89]. Full entries are listed in the References section of the CPMN book.