17.17 Burns in Children*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • Crying in pain **Objective** • Scalds over 15% TBSA • High behavioral pain score	**ACTUAL** Acute pain related to burn injury as evidenced by crying and an elevated pain score.	• Comfortable at rest and during dressing changes.	• Give weight-based analgesia before dressings. • Use play to distract him. • Keep the burns covered.	• Achieved when he is comfortable.
Subjective • Mother says, “He has not passed much urine.” **Objective** • Low urine output • Tachycardia	**ACTUAL** Inadequate fluid volume related to fluid loss through burns as evidenced by low urine output.	• Urine 1 mL/kg/hour within 8 hours.	• Give fluids by the prescribed burn formula. • Measure urine hourly.	• Outcome reached when output is on target.
Subjective *No specific complaint* **Objective** • Open burn wounds	**RISK** Risk for infection as evidenced by risk factors: loss of skin barrier.	• Burns free of infection.	• Use aseptic technique for dressings. • Watch temperature and the wounds. • Give tetanus prophylaxis as ordered.	• Considered met if no infection develops.

* **References for this care plan:** [35], [38], [39], [49]. Full entries are listed in the References section of the CPMN book.