2.17 Total Parenteral Nutrition*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • Lost 6 kg in a month **Objective** • Underweight • Low albumin • Cannot tolerate enteral feeds	**ACTUAL** Inadequate nutritional intake related to inability to absorb nutrients as evidenced by weight loss and hypoalbuminemia.	• Weight steady or rising within 1 week.	• Run TPN by pump at the prescribed rate. • Never stop TPN abruptly; taper as ordered. • Weigh daily at a fixed time. • Follow electrolytes, glucose, liver function and triglycerides. • Reserve the TPN lumen for TPN alone.	• Met if weight is steady or increasing.
Subjective • "My mouth feels dry." **Objective** • Dry oral mucosa • Nil by mouth	**ACTUAL** Impaired oral mucous membrane integrity related to nil-by-mouth status as evidenced by dry oral mucosa.	• Moist mucosa within 24 hours.	• Give mouth care every 4 hours with a soft brush. • Moisturize the lips. • Use the prescribed mouth rinse. • Look daily for ulcers or thrush.	• Achieved when the mucosa is moist.
Subjective *No specific complaint* **Objective** • High-dextrose infusion • Critically ill	**RISK** Risk for physical injury as evidenced by risk factors: hyperglycemia and hypoglycemia associated with high-dextrose parenteral nutrition.	• Blood glucose stays within target.	• Check glucose every 6 hours, or as ordered. • Give insulin as prescribed. • Report symptoms of hypo- or hyperglycemia.	• Outcome reached while glucose remains in range.

* **References for this care plan:** [35], [37], [55], [105], [108]. Full entries are listed in the References section of the CPMN book.