2.7 Pressure Injury Prevention*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "It is hard for me to change position." **Objective** • Cannot turn without help • Lies in one position for long periods	**ACTUAL** Impaired bed mobility related to weakness as evidenced by inability to reposition independently.	• Repositioned at least two-hourly with no lasting redness.	• Follow a two-hourly turning chart. • Pad between bony prominences with pillows. • Use slide sheets to avoid friction and shear. • Encourage him to help with turns where he can.	• Achieved: skin intact, no non-blanching redness.
Subjective • "I cannot reach the toilet in time." **Objective** • Wet bed linen • Unable to walk to the toilet alone	**ACTUAL** Disability-associated urinary incontinence related to impaired mobility as evidenced by urine leakage before reaching the toilet.	• Skin stays dry and incontinent episodes fall within 3 days.	• Offer a bedpan or urinal every 2 hours. • Clean and dry the skin promptly after each episode. • Apply barrier cream. • Avoid layers of pads and sheets beneath him.	• Achieved when his skin is dry and intact.
Subjective • Says he has no appetite **Objective** • Braden score 12 • Thin, with prominent bony areas • Low serum albumin	**RISK** Risk for adult pressure injury as evidenced by risk factors: immobility, moisture, inadequate nutrition and a low Braden score.	• No pressure injury develops during the stay.	• Inspect skin over bony prominences each shift. • Nurse him on a pressure-redistributing mattress. • Keep the head of the bed at 30 degrees or less unless contraindicated. • Refer to the dietitian and encourage protein intake. • Repeat the Braden score daily.	• Outcome reached if no pressure injury forms. • Non-blanching redness is documented and reported.

*References for this care plan: [6], [19], [35], [68], [110]. Full entries are listed in the References section.