3.15 Tracheostomy Care*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • Points to his throat to show discomfort **Objective** • Thick secretions in the tracheostomy tube • Noisy breathing	**ACTUAL** Ineffective airway clearance related to increased secretions and an artificial airway as evidenced by noisy breathing and visible secretions.	• Tracheostomy patent and airway clear.	• Suction when indicated, using sterile technique. • Clean or change the inner cannula per policy. • Humidify the inspired air. • Keep a spare tube, obturator and tracheal dilator at the bedside.	• Outcome reached while the airway is clear.
Subjective • "I do not want people to see my neck." **Objective** • Will not look at the stoma • Covers his neck when visitors arrive	**ACTUAL** Disrupted body image related to tracheostomy as evidenced by avoidance of looking at the stoma.	• Looks at the stoma and starts to share in its care.	• Encourage him to talk about how he feels. • Involve him in care a step at a time. • Suggest a scarf or stoma cover when he goes out.	• Considered met when he takes part in stoma care.
Subjective *No specific complaint* **Objective** • Surgical stoma • Secretions around the stoma	**RISK** Risk for infection as evidenced by risk factors: open stoma and secretions.	• Stoma clean, with no sign of infection.	• Clean the stoma and change the dressing per policy. • Change soiled ties, leaving room for one finger. • Look for redness or pus.	• Goal reached while the stoma stays clean.

* **References for this care plan:** [35], [37], [110], [128]. Full entries are listed in the References section of the CPMN book.