4.3 Angina Pectoris*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I get chest pressure when I walk fast; it goes when I rest." **Objective** • Episodes last 5 minutes • Relieved by nitroglycerin • ST depression on ECG during pain	**ACTUAL** Acute pain related to myocardial ischemia as evidenced by exertional chest pressure relieved by rest.	• Chest pain relieved within 5–15 minutes of treatment.	• Stop the activity; sit or lie him down. • Give sublingual nitroglycerin as prescribed, checking BP before each dose. • Give oxygen if SpO_2 is below target. • Record a 12-lead ECG while pain is present. • Report pain that persists after three doses, as this may be an MI.	• Outcome reached when pain is relieved.
Subjective • "I do not know how much activity is safe." **Objective** • Newly diagnosed • Cannot describe how to use nitroglycerin	**ACTUAL** Inadequate health knowledge related to a new diagnosis as evidenced by questions about activity and medication.	• Explains how to use nitroglycerin and when to get help.	• Teach him to carry nitroglycerin, keep it in a dark container and replace it as directed. • Tell him to call emergency services if pain continues after the first dose, following local advice. • Discuss building up exercise gradually and avoiding heavy meals before exertion.	• Considered met when he explains correct use.
Subjective *No specific complaint* **Objective** • Coronary artery disease • Smoker • High cholesterol	**RISK** Risk for impaired cardiovascular function as evidenced by risk factors: coronary artery disease, smoking and hyperlipidemia.	• Commits to reducing his risk factors.	• Support smoking cessation. • Teach a diet low in saturated fat. • Encourage regular statin and antiplatelet use.	• Goal reached when he describes a risk-reduction plan.

*References for this care plan: [11], [17], [32], [35]. Full entries are listed in the References section of the CPMN book.