10.13 Anaphylaxis

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "My throat is closing." **Objective** • Stridor and wheeze • Face and lips swollen • Hives	**ACTUAL** **Ineffective airway clearance** related to airway edema and bronchospasm as evidenced by stridor and wheeze.	Airway open within minutes of treatment.	• Call for help and remove the trigger. • Give IM adrenaline straight away, following the protocol. • Give high-flow oxygen. • If the swelling gets worse, prepare for an advanced airway. • No improvement? • Repeat the adrenaline as the protocol says.	Airway open: **goal met**.
Subjective • "I feel faint." **Objective** • BP 80/40 mmHg • HR 130/min	**ACTUAL** **Decreased cardiac output** related to vasodilation and capillary leak as evidenced by hypotension and tachycardia.	Systolic BP above 90 mmHg within 30 minutes.	• Lay him flat with his legs up (sit him up if lying flat makes the breathing worse). • Give an IV fluid bolus as prescribed. • Continuous monitoring of BP, HR and SpO_2. • After the adrenaline, give the antihistamine and steroid as prescribed.	BP back up: **goal met**.
Subjective • *No specific complaint* **Objective** • Has a known severe allergy • May have a second (biphasic) reaction	**RISK** **Risk for allergy reaction** as evidenced by risk factors: known severe allergy.	Avoids exposure in future and uses the auto-injector correctly.	• Once he recovers, observe him for the time set by policy. • Write the allergy clearly in the notes and on his wristband. • Teach him how to use the adrenaline auto-injector. • Refer to the allergy clinic.	Shows you correctly: **goal met**.