10.2 Sickle Cell Crisis

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "The pain in my bones is unbearable." • Gives the pain 10/10 **Objective** • Known to have sickle cell disease • Guards his limbs • HR 120/min	**ACTUAL** **Acute pain** related to vaso-occlusion and tissue ischemia as evidenced by severe bone pain.	Within 1 hour of analgesia, says the pain is bearable.	• Give strong analgesia without delay, as prescribed (often opioids by protocol). • Check pain again every 30–60 minutes. • Put warmth on the painful areas; keep cold away. • Believe him when he tells you about his pain.	Pain bearable: **goal met**.
Subjective • "I am thirsty." **Objective** • Mouth dry • Urine concentrated	**ACTUAL** **Deficient fluid volume** related to inadequate intake and increased loss as evidenced by dry mucosa.	Well hydrated within 24 hours.	• Give IV and oral fluids as prescribed. • Keep an I&O chart. • If SpO_2 is below target, give oxygen.	Hydrated: **goal met**.
Subjective • "I have chest pain and cough." **Objective** • In sickle crisis • Breathing shallow because of pain	**RISK** **Risk for infection** as evidenced by risk factors: functional asplenia and lung involvement (acute chest syndrome).	Chest syndrome and infection picked up early.	• Watch SpO_2, RR and temperature. • Encourage the incentive spirometer. • Report chest pain, fever or a drop in SpO_2 at once.	No complications: **goal met**.