10.7 Disseminated Intravascular Coagulation (DIC)

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • *No specific complaint* **Objective** • Oozing from IV and puncture sites • Blood in the urine • Platelets and fibrinogen low, D-dimer high • BP 92/58 mmHg	**ACTUAL** **Deficient fluid volume** related to active bleeding from coagulopathy as evidenced by oozing, hematuria and falling BP.	Within 24 hours, bleeding less and vital signs stable.	• Help treat whatever caused it (sepsis, an obstetric complication, trauma). • Give platelets, plasma and cryoprecipitate as prescribed. • Press on puncture sites for longer than usual. • Follow vital signs, Hb and clotting results. • Avoid any puncture that is not needed.	Less bleeding, stable: **goal met**.
Subjective • *No specific complaint* **Objective** • Fingers and toes cool and mottled • Capillary refill slow	**ACTUAL** **Ineffective peripheral tissue perfusion** related to microvascular thrombi as evidenced by mottled, cool extremities.	Hands and feet warm, with capillary refill under 3 seconds.	• Check hands and feet every hour. • Keep her warm. • Report cyanosis or any new organ failure.	Perfusion better: **goal met**.
Subjective • *No specific complaint* **Objective** • Clotting problem continuing • Septic	**RISK** **Risk for shock** as evidenced by risk factors: hemorrhage and sepsis.	MAP above 65 mmHg and urine output at least 0.5 mL/kg/hour.	• Follow hemodynamics and hourly urine output. • Give fluids and blood products as ordered.	Stable: **goal met**.