12.4 Tonsillectomy (Adult)

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "It hurts a lot to swallow." • Gives the pain 7/10 **Objective** • Throat pain • Does not want to eat	**ACTUAL** **Acute pain** related to surgical trauma of the throat as evidenced by pain on swallowing.	Pain low enough to eat and drink.	• Give analgesia regularly, especially before meals. • Encourage cool drinks and soft food. • Encourage a normal diet early, because it helps healing.	Eating and drinking: **goal met**.
Subjective • "I am not drinking much because of pain." **Objective** • Drinking less • Mouth dry	**ACTUAL** **Deficient fluid volume** related to reduced intake from pain as evidenced by dry mucosa.	Within 24 hours, drinking at least 1.5 L a day.	• Chart what he drinks. • Offer drinks often. • IV fluids if ordered.	Intake adequate: **goal met**.
Subjective • *No specific complaint* **Objective** • Tonsil bed	**RISK** **Risk for bleeding** as evidenced by risk factors: surgical wound in a vascular area.	No bleeding after surgery.	• Watch for frequent swallowing, fresh blood and a rising heart rate. • Tell him to report any bleeding after he goes home (especially around day 5–10).	No bleeding: **goal met**.