13.5 Prostate Cancer and Prostatectomy

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I leak urine since the catheter was removed." **Objective** • Dribbling since the catheter came out • Uses 3 pads a day	**ACTUAL** **Stress urinary incontinence** related to sphincter weakness after prostatectomy as evidenced by urine leakage.	Within 6–12 weeks, uses fewer pads.	• Teach pelvic floor exercises and check he is doing them right. • Give him suitable continence products. • He should cut down caffeine and alcohol. • Explain that continence often gets better over the months.	Continence improving: **goal met**.
Subjective • "Will I be able to have relations with my wife?" **Objective** • Worried about erectile function after surgery • Avoids talking about it with his wife	**ACTUAL** **Sexual dysfunction** related to effects of radical prostatectomy as evidenced by expressed concern about erectile function.	Talks about his worries and the options open to him.	• Give him privacy and do not judge. • Give accurate information about recovery and the treatments available. • If he wishes, include his wife. • Refer to urology or a sexual health specialist.	Discusses the options: **goal met**.
Subjective • *No specific complaint* **Objective** • Had pelvic surgery • Has cancer	**RISK** **Risk for thrombosis** as evidenced by risk factors: pelvic surgery and cancer.	No DVT.	• Give the prophylactic anticoagulant as ordered. • Get him moving early. • Teach leg exercises and how to check his calves.	No DVT: **goal met**.