14.10 Poisoning and Drug Overdose

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • Family found him with empty tablet bottles **Objective** • RR 8/min and shallow • SpO_2 86% • Pupils pinpoint	**ACTUAL** **Ineffective breathing pattern** related to central nervous system depression as evidenced by bradypnea and hypoxemia.	RR 12–20/min and SpO_2 94% or more.	• Open the airway and support breathing with a bag-valve-mask if needed. • Give the antidote as prescribed (naloxone for opioids, for example). • Continuous monitoring of RR, SpO_2 and level of consciousness. • Contact the poison information service as policy says. • Take blood for drug levels as ordered.	Breathing adequate: **goal met**.
Subjective • *No specific complaint* **Objective** • GCS 10 • Disoriented when he wakes	**ACTUAL** **Acute confusion** related to toxic effects of substance as evidenced by reduced GCS and disorientation.	Back to his usual orientation.	• Check GCS often. • Keep him safe and under close observation. • Once he is alert, ask whether the overdose was intentional and refer to the mental health team.	Oriented and referred: **goal met**.
Subjective • *No specific complaint* **Objective** • Not fully conscious • Vomiting	**RISK** **Risk for aspiration** as evidenced by risk factors: decreased level of consciousness and vomiting.	Airway stays clear, with no aspiration.	• Put him in the recovery position. • Have suction ready. • Nothing by mouth until he is alert.	No aspiration: **goal met**.