# 14.11 Heat Stroke

| Assessment | Nursing Diagnosis | Plan / Expected Outcome | Nursing Interventions | Evaluation |
|---|---|---|---|---|
| **Subjective**<br>• Collapsed while working outside in summer<br>**Objective**<br>• Core temperature 41 °C<br>• Skin hot and dry<br>• Confused | **ACTUAL**<br>**Hyperthermia** related to exposure to high environmental temperature as evidenced by core temperature 41 °C, hot skin and confusion. | Core temperature below 39 °C within 30 minutes. | • Get him somewhere cool and take off his clothes.<br>• Cool him fast: cold water immersion, or evaporative cooling with fans plus ice packs to the neck, armpits and groin.<br>• Monitor core temperature continuously and stop active cooling at about 39 °C, as the protocol says.<br>• Watch for seizures. | Temperature on target: **goal met**. |
| **Subjective**<br>• *No specific complaint*<br>**Objective**<br>• HR 130/min<br>• BP 90/60 mmHg<br>• Passing small amounts of dark urine | **ACTUAL**<br>**Deficient fluid volume** related to excessive sweating and heat exposure as evidenced by tachycardia, hypotension and low urine output. | Urine output adequate and vital signs stable. | • Give IV fluids as prescribed.<br>• Record urine output every hour and look for dark urine (rhabdomyolysis).<br>• Follow electrolytes, creatinine and CK. | Stable: **goal met**. |
| **Subjective**<br>• *No specific complaint*<br>**Objective**<br>• Brain affected by the heat<br>• Confused | **RISK**<br>**Risk for injury** as evidenced by risk factors: seizures and confusion from heat stroke. | No injury. | • Seizure precautions in place.<br>• If he has a seizure, give the anticonvulsant as prescribed.<br>• Keep him under close observation. | No injury: **goal met**. |