14.2 Sepsis and Septic Shock

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I feel very unwell and shivery." **Objective** • Temperature 39.3 °C or below 36 °C • BP 85/45 mmHg, HR 128/min, RR 28/min • Lactate 4 mmol/L • Newly confused	**ACTUAL** **Ineffective peripheral tissue perfusion** related to vasodilation and capillary leak as evidenced by hypotension, raised lactate, mottled skin and confusion.	Within 6 hours, MAP 65 mmHg or more and lactate falling.	• Follow the sepsis bundle: blood cultures, then IV antibiotics within 1 hour. • Give an IV fluid bolus as prescribed, then reassess. • Measure lactate and repeat it as ordered. • If BP stays low, start vasopressors as prescribed. • Record urine output every hour.	MAP on target, lactate falling: **goal met**.
Subjective • *No specific complaint* **Objective** • Temperature 39.3 °C • Shaking with rigors	**ACTUAL** **Hyperthermia** related to infection as evidenced by temperature 39.3 °C and rigors.	Temperature below 38 °C within 6 hours.	• Give the antipyretic as prescribed. • Take off extra covers. • Temperature every hour.	Temperature coming down: **goal met**.
Subjective • *No specific complaint* **Objective** • In septic shock • BP low	**RISK** **Risk for decreased cardiac output** as evidenced by risk factors: septic cardiomyopathy and dysrhythmias.	Rhythm and output stable.	• Continuous ECG monitoring. • Follow electrolytes. • Report any new dysrhythmia.	Stable: **goal met**.