14.6 Multiple Trauma

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I feel weak and dizzy." **Objective** • Fractured pelvis and femur • BP 86/54 mmHg, HR 128/min • Skin pale and cold	**ACTUAL** **Deficient fluid volume** related to hemorrhage from multiple injuries as evidenced by hypotension, tachycardia and cold skin.	MAP 65 mmHg or more and urine output at least 0.5 mL/kg/hour.	• Do the primary survey: airway with cervical spine control, breathing, circulation. • Stop external bleeding and put on a pelvic binder as ordered. • Give warmed blood products as prescribed (massive transfusion protocol). • Vital signs every 5–15 minutes and urine output every hour. • Get him ready for imaging and surgery.	Hemodynamics stable: **goal met**.
Subjective • "Everything hurts." • Gives the pain 9/10 **Objective** • Several fractures • Grimacing • Guarding	**ACTUAL** **Acute pain** related to multiple tissue injuries as evidenced by pain score 9/10 and guarding.	Within 1 hour, says the pain is bearable.	• Give IV analgesia in small doses, titrated, as prescribed. • Splint the fractures. • Check pain and sedation often. • Handle and move him gently.	Pain bearable: **goal met**.
Subjective • *No specific complaint* **Objective** • Open wounds • Has invasive lines and drains	**RISK** **Risk for infection** as evidenced by risk factors: open wounds, contamination and invasive devices.	No wound infection.	• Clean and dress wounds with aseptic technique. • Give tetanus prophylaxis and antibiotics as ordered. • Watch his temperature and how the wounds look.	No infection: **goal met**.