15.12 Preterm Labor

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I have regular tightening at 30 weeks." **Objective** • Contracting every 5 minutes • Cervix opening	**ACTUAL** **Labor pain** related to preterm uterine contractions as evidenced by regular contractions.	Copes with the contractions while treatment is given.	• Give tocolytics as prescribed. • Use comfort measures and lay her on her left side. • Time how often the contractions come.	Controlled: **goal met**.
Subjective • "It is too early for my baby." **Objective** • Anxious • Tearful	**ACTUAL** **Anxiety** related to threat of preterm birth as evidenced by worry.	Says she feels less anxious.	• Explain the plan of care. • Arrange for the neonatal team to visit her. • Let a support person stay.	Calmer: **goal met**.
Subjective • *No specific complaint* **Objective** • Preterm • Cervix opening	**RISK** **Risk for disturbed maternal-fetal dyad** as evidenced by risk factors: preterm labor.	Everything is ready for the baby's preterm birth.	• Give corticosteroids as prescribed. • Give magnesium sulfate for neuroprotection as prescribed. • Monitor the fetal heart. • Let the neonatal unit know.	Ready: **goal met**.