15.17 Normal Postpartum Care

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "My stitches hurt when I sit." • Gives the pain 5/10 **Objective** • Perineal tear repaired A little swollen	**ACTUAL** **Acute pain** related to perineal trauma as evidenced by pain on sitting and swelling.	Pain 3/10 or less.	• In the first 24 hours, put a cold pack on the perineum. • Give analgesia as prescribed. • Teach perineal hygiene and to change pads often.	Pain controlled: **goal met**.
Subjective • "How do I care for the baby's cord and bath?" **Objective** • First baby • Not sure how to bathe the baby	**ACTUAL** **Deficient knowledge** related to newborn care and self-care as evidenced by questions.	Before discharge, shows you basic baby care.	• Teach cord care, bathing, feeding cues and safe sleep. • Teach the danger signs in a newborn: feeding poorly, jaundice, fever, difficulty breathing. • Teach her own warning signs: heavy bleeding, fever, pain in the leg, low mood.	Shows you: **goal met**.
Subjective • *No specific complaint* **Objective** • Just given birth	**RISK** **Risk for bleeding** as evidenced by risk factors: postpartum uterus.	Fundus firm, lochia normal.	• In the first 24 hours, check the fundus and lochia regularly. • Encourage her to pass urine (a full bladder stops the uterus contracting). • Teach her fundal massage.	No hemorrhage: **goal met**.