15.18 Postpartum Hemorrhage

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I feel faint and wet." **Objective** • Has lost more than 500 mL of blood • Uterus soft and boggy • HR 120/min, BP 90/55 mmHg	**ACTUAL** **Deficient fluid volume** related to excessive blood loss from uterine atony as evidenced by heavy bleeding, soft uterus, tachycardia and hypotension.	Within 1 hour, bleeding controlled and vital signs stable.	• Call for help. • Rub up the uterus and empty the bladder. • Give uterotonics as prescribed. • Put in two large-bore IV cannulas and give fluids and blood as ordered. • Measure the blood loss and check vital signs every 5–15 minutes.	Bleeding controlled: **goal met**.
Subjective • "Am I going to die?" **Objective** • Frightened • Crying	**ACTUAL** **Fear** related to heavy bleeding and emergency interventions as evidenced by expressed fear.	Says she feels safer as the bleeding stops.	• Stay with her. • Explain what you are doing in a few words. • Keep her husband informed.	Calmer: **goal met**.
Subjective • *No specific complaint* **Objective** • Still losing blood	**RISK** **Risk for shock** as evidenced by risk factors: hemorrhage.	MAP 65 mmHg or more and urine output adequate.	• Put in a urinary catheter and record the output. • Check clotting. • If the bleeding goes on, get her ready for theatre.	Stable: **goal met**.