15.22 Breastfeeding Support

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "My baby does not latch and keeps crying." **Objective** • Latch shallow • Clicking sound while feeding • Baby unsettled after feeds	**ACTUAL** **Ineffective breastfeeding** related to poor latch technique as evidenced by shallow latch and unsettled infant.	Within 48 hours, latching well and the baby is satisfied.	• Watch a whole feed. • Teach positioning: baby's body held close, nose to nipple, mouth wide open. • Encourage feeding on demand, 8–12 times in 24 hours. • Teach the signs of a good feed: wet diapers, hearing the baby swallow.	Feeding well: **goal met**.
Subjective • "My nipples are sore." **Objective** • Nipples red • Winces when the baby latches	**ACTUAL** **Acute pain** related to nipple trauma from poor latch as evidenced by sore nipples.	Within 48 hours, says her nipples are less sore.	• Correct the latch. • After feeds, rub a little expressed breast milk on the nipples. • Change feeding positions.	Less pain: **goal met**.
Subjective • *No specific complaint* **Objective** • Latch poor	**RISK** **Risk for impaired skin integrity** as evidenced by risk factors: repeated nipple trauma.	Nipple skin stays intact.	• Check the latch at every feed. • Look at the nipples every day. • Let the nipples air-dry after feeds.	Intact: **goal met**.