15.7 Gestational Hypertension and Pre-eclampsia

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "My face and hands are swollen." **Objective** • BP 156/102 mmHg • Protein in the urine • Face and hands swollen	**ACTUAL** **Excess fluid volume** related to reduced renal perfusion and capillary leak as evidenced by edema and proteinuria.	BP on the prescribed target and edema not getting worse.	• BP every 4 hours, or as ordered. • Test urine for protein and keep an I&O chart. • Give antihypertensives as prescribed. • Follow liver enzymes, platelets and creatinine. • Keep checking on the baby.	BP controlled: **goal met**.
Subjective • "I am afraid for my baby." **Objective** • Anxious • Keeps asking about the baby	**ACTUAL** **Anxiety** related to threat to self and fetus as evidenced by worry.	Says she feels less anxious and can name the warning signs.	• Explain the plan of care and the monitoring. • Teach the warning signs: severe headache, problems with vision, pain in the upper abdomen, baby moving less. • Let a support person stay.	Calmer: **goal met**.
Subjective • *No specific complaint* **Objective** • Pre-eclampsia is severe	**RISK** **Risk for injury** as evidenced by risk factors: possible eclamptic seizure.	No seizures.	• Give magnesium sulfate as prescribed. • While she is on magnesium, check reflexes, RR and urine output. • Keep calcium gluconate at hand. • Keep the room quiet and dim.	No seizure: **goal met**.