15.8 Eclampsia

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • *No specific complaint* **Objective** • Generalised seizure • BP 170/110 mmHg • Frothy secretions	**ACTUAL** **Ineffective airway clearance** related to seizure activity as evidenced by frothy secretions.	Airway clear and seizure controlled.	• Call for help and turn her onto her left side. • Keep the airway open; suction and give oxygen. • Give magnesium sulfate as prescribed. • Do not hold her down; protect her from injury.	Seizure controlled: **goal met**.
Subjective • Her husband says • "She does not know where she is." **Objective** • Drowsy after the seizure	**ACTUAL** **Acute confusion** related to postictal state as evidenced by drowsiness and disorientation.	Oriented within a few hours of the seizure.	• Neuro checks every 15–30 minutes. • Reorient her gently. • Keep her under close observation with the side rails up.	Oriented: **goal met**.
Subjective • *No specific complaint* **Objective** • Mother has had a seizure	**RISK** **Risk for disturbed maternal-fetal dyad** as evidenced by risk factors: maternal hypoxia during seizures.	Baby stays well until it can be delivered safely.	• Continuous fetal heart monitoring. • Give the mother oxygen. • Once the mother is stable, get her ready for delivery.	Safe outcome: **goal met**.