16.6 Birth Asphyxia

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • *No specific complaint* **Objective** • Apgar 3 at 1 minute • Breathing weak	**ACTUAL** **Impaired spontaneous ventilation** related to hypoxic insult at birth as evidenced by poor respiratory effort.	Starts breathing effectively on its own.	• Start neonatal resuscitation according to protocol. • Watch HR and SpO_2. • Move the baby to the neonatal unit for ongoing support.	Breathing on its own: **goal met**.
Subjective • *No specific complaint* **Objective** • Seizures • Tone abnormal	**ACTUAL** **Decreased intracranial adaptive capacity** related to hypoxic brain injury as evidenced by seizures and abnormal tone.	Seizures controlled.	• Give anticonvulsants as prescribed. • If the baby is eligible, start therapeutic cooling according to protocol. • Watch neurological status and blood glucose.	Controlled: **goal met**.
Subjective • *No specific complaint* **Objective** • Baby in the neonatal unit • Parents afraid to touch the baby	**RISK** **Risk for impaired attachment** as evidenced by risk factors: separation of parents and infant.	Parents visit and join in with care.	• Encourage them to visit and to touch the baby. • Keep the parents informed. • Let them do simple care tasks.	Parents involved: **goal met**.