17.1 Bronchiolitis

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • Mother says • "He is breathing fast and not feeding." **Objective** • 4-month-old, RR 64/min • Nasal flaring, subcostal recession • Wheeze and crackles, SpO_2 90%	**ACTUAL** **Ineffective airway clearance** related to excessive mucus and small airway inflammation as evidenced by retractions, crackles and low SpO_2.	Within 48 hours, SpO_2 92% or more (or the local target) and less work of breathing.	• Before feeds, clear the nose with saline drops and gentle suction. • Give oxygen to keep SpO_2 on target. • Lay him with his head slightly raised. • Watch RR, SpO_2 and any sign of exhaustion or apnea.	SpO_2 on target, less distressed: **goal met**.
Subjective • Mother says • "He takes only a little milk." **Objective** • Taking less than half his usual feeds • Fewer wet diapers than usual	**ACTUAL** **Ineffective infant feeding dynamics** related to breathlessness during feeds as evidenced by reduced intake and fewer wet diapers.	Stays well hydrated.	• Offer small feeds, often. • If he is not taking enough, give NG feeds or IV fluids as ordered. • Count wet diapers and weigh every day.	Hydrated: **goal met**.
Subjective • *No specific complaint* **Objective** • Young baby • Chest infection	**RISK** **Risk for infection (to others)** as evidenced by risk factors: contagious viral infection.	Infection not passed to other babies.	• Contact and droplet precautions as policy says. • Strict hand hygiene. • Teach the parents hand hygiene.	No cross-infection: **goal met**.