17.19 Pyloric Stenosis

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • Mother says • "He vomits forcefully after every feed." **Objective** • Projectile vomiting, not bile-stained • Olive-shaped mass felt • Fontanelle sunken, few wet diapers	**ACTUAL** **Deficient fluid volume** related to persistent vomiting as evidenced by sunken fontanelle and reduced wet diapers.	Rehydrated, with normal electrolytes, before surgery.	• Nothing by mouth; give IV fluids as prescribed. • Follow electrolytes and acid–base status. • Record vomiting, I&O and weight.	Rehydrated: **goal met**.
Subjective • *No specific complaint* **Objective** • Losing weight • Hungry straight after vomiting	**ACTUAL** **Imbalanced nutrition: less than body requirements** related to vomiting as evidenced by weight loss.	Tolerates feeds after pyloromyotomy.	• After surgery, bring feeds back as ordered. • Burp him often. • Weigh every day.	Tolerating feeds: **goal met**.
Subjective • *No specific complaint* **Objective** • Losing gastric acid	**RISK** **Risk for electrolyte imbalance** as evidenced by risk factors: vomiting of gastric contents (metabolic alkalosis, hypokalemia).	Electrolytes and acid–base balance back to normal before surgery.	• Follow electrolytes. • Replace as prescribed.	Normal: **goal met**.