# 17.22 Hypospadias Repair

| Assessment | Nursing Diagnosis | Plan / Expected Outcome | Nursing Interventions | Evaluation |
|---|---|---|---|---|
| **Subjective**<br>• Crying after the operation<br>**Objective**<br>• Dressing on the penis<br>• Urinary stent in place<br>• Crying | **ACTUAL**<br>**Acute pain** related to surgery as evidenced by crying. | Looks comfortable within 1 hour of analgesia. | • Give analgesia by weight as prescribed.<br>• Give the medicine for bladder spasm as ordered.<br>• Comfort and distract him. | Comfortable: **goal met**. |
| **Subjective**<br>• Mother asks<br>• "How do I care for the tube at home?"<br>**Objective**<br>• Stent in situ<br>• Mother not sure how to put the diaper on | **ACTUAL**<br>**Deficient knowledge (parent)** related to stent and dressing care as evidenced by questions. | Parent shows you how she will care for it. | • Teach double-diapering, keeping the stent free.<br>• Teach the signs of infection. | Shows you: **goal met**. |
| **Subjective**<br>• *No specific complaint*<br>**Objective**<br>• Surgical site<br>• Urinary stent in place | **RISK**<br>**Risk for infection** as evidenced by risk factors: surgical wound and urinary stent. | Surgical site stays free of infection. | • Keep the area clean.<br>• Give antibiotics as ordered.<br>• Teach the parents the signs of infection. | No infection: **goal met**. |