19.11 Incontinence in Older Adults at Home

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I cannot reach the toilet in time at night." **Objective** • Urge incontinence • Up twice a night to pass urine	**ACTUAL** **Urge urinary incontinence** related to overactive bladder as evidenced by leakage before reaching the toilet.	Within 4 weeks, fewer episodes.	• Keep a bladder diary. • Bladder training with voiding at set times. • Cut down caffeine. • A commode by the bed at night.	Fewer episodes: **goal met**.
Subjective • "I am ashamed." **Objective** • Avoids visitors • Uses a lot of pads	**ACTUAL** **Situational low self-esteem** related to incontinence as evidenced by shame and avoiding visitors.	Talks about how she feels and accepts help.	• Talk about discreet continence products. • Talk with her supportively. • Reassure her that treatment can help.	Accepts help: **goal met**.
Subjective • *No specific complaint* **Objective** • Skin damp • Perineum red	**RISK** **Risk for impaired skin integrity** as evidenced by risk factors: moisture.	Skin stays intact.	• Teach skin care and the use of barrier cream. • Change pads quickly. • Look at the skin every day.	Intact: **goal met**.