2.10 Deficient Fluid Volume

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I feel thirsty and dizzy." **Objective** • BP 95/60 mmHg, HR 115/min • Urine output 20 mL/hour, urine dark • Skin turgor poor	**ACTUAL** **Deficient fluid volume** related to active fluid loss as evidenced by hypotension, tachycardia and low urine output.	Within 8 hours, urine output is at least 0.5 mL/kg/hour and vital signs are stable.	• Vital signs every hour until stable. • Give IV fluids as prescribed and keep checking the rate. • Record urine output every hour, along with I&O. • Weigh daily, at the same time each day. • Follow serum electrolytes, urea and creatinine. • Once allowed, encourage drinking.	BP and HR normal, urine output adequate: **goal met**. If not, ask the physician to review the fluid plan.
Subjective • "My mouth is very dry." **Objective** • Lips dry and cracked • Dry tongue	**ACTUAL** **Impaired oral mucous membrane integrity** related to dehydration as evidenced by dry, cracked lips and tongue.	Mouth moist again within 24 hours.	• Mouth care every 4 hours. • Put moisturiser on the lips. • If allowed, give small sips of water often.	Mouth moist: **goal met**.
Subjective • Still vomiting **Objective** • Fluid loss continuing • BP dropping	**RISK** **Risk for shock** as evidenced by risk factors: hypovolemia.	MAP stays above 65 mmHg and level of consciousness stays normal.	• Check level of consciousness and capillary refill. • Report a systolic BP below 90 mmHg. • Keep a large-bore IV line in place. • Be ready for fluid resuscitation as ordered.	No signs of shock: **goal met**.