2.14 Blood Transfusion

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I get tired after walking a few steps." **Objective** • Hb 6.8 g/dL • Pale • HR 110/min on exertion	**ACTUAL** **Activity intolerance** related to reduced oxygen-carrying capacity as evidenced by fatigue on exertion and low Hb.	After the transfusion, manages self-care without becoming breathless.	• Plan rest between activities. • Give blood as prescribed. • Check HR and SpO_2 while she is active. • Recheck Hb after the transfusion, as ordered.	Does her ADL without tiring: **goal met**.
Subjective • "I am worried about receiving blood." **Objective** • Face looks tense • Keeps asking if it is safe	**ACTUAL** **Anxiety** related to unfamiliar procedure as evidenced by worry and repeated questions.	Before the transfusion starts, she says she understands and feels less anxious.	• Explain what will happen, how long it takes and what checks are done. • Tell her which symptoms she must report. • Stay with her for the first 15 minutes.	Calm and cooperative: **goal met**.
Subjective • *No specific complaint* **Objective** • Receiving a blood product • Has had transfusions before	**RISK** **Risk for injury** as evidenced by risk factors: possible transfusion reaction or incompatibility.	Transfusion finished with no reaction.	• At the bedside, check the patient's identity and the blood unit with a second nurse. • Take baseline vital signs. • Run it slowly at first and watch closely for the first 15 minutes. • Vital signs as per policy. • If there is a reaction: stop the transfusion, keep the line open with normal saline, call the physician and send the unit back. • Finish it within 4 hours of the unit leaving storage.	No reaction: **goal met**.