2.23 Use of Physical Restraints

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • Family says he does not know where he is **Objective** • Disoriented • Pulls at his lines and tubes	**ACTUAL** **Acute confusion** related to (cause, for example delirium) as evidenced by disorientation and pulling at devices.	Kept safe using the least restrictive measures.	• Try other options first: reorientation, a family member at the bedside, close observation. • Use a restraint only with a physician's order and as policy allows. • Explain to the family what is happening and why.	Safe, and restraint off as soon as possible: **goal met**.
Subjective • Asks to be able to move his arms **Objective** • Wrist restraints on Can hardly move his arms	**ACTUAL** **Impaired physical mobility** related to restraint use as evidenced by limited movement.	Circulation and joint movement kept normal.	• Every 2 hours, release the restraint and put the joints through ROM. • Every hour, check circulation, skin and sensation. • Offer drinks and the toilet.	Circulation and skin normal: **goal met**.
Subjective • *No specific complaint* **Objective** • Restraint on Confusion	**RISK** **Risk for injury** as evidenced by risk factors: physical restraint and agitation.	No injury.	• Use the right size of restraint and tie it to the bed frame, never the side rails. • Write down why it is on, every check made and how he responds. • Keep asking whether it is still needed.	No injury: **goal met**.