2.7 Pressure Injury Prevention

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "It is hard for me to change position." **Objective** • Cannot turn without help • Lies in the same position for long periods	**ACTUAL** **Impaired bed mobility** related to weakness as evidenced by inability to reposition independently.	Repositioned at least every 2 hours, with no pressure redness.	• Turn every 2 hours, following a turning chart. • Put pillows between bony prominences so they do not touch. • Use slide sheets so the skin is not dragged. • Ask the patient to help with turning where he can.	Skin intact, no redness that stays: **goal met**.
Subjective • "I cannot reach the toilet in time." **Objective** • Bed linen wet • Cannot walk to the toilet alone	**ACTUAL** **Functional urinary incontinence** related to impaired mobility as evidenced by urine leakage before reaching toilet.	Within 3 days his skin is dry and he is incontinent less often.	• Offer a bedpan or urinal every 2 hours. • Clean and dry the skin quickly after every episode. • Put on barrier cream. • Do not pile layers of pads and sheets under him.	Skin dry and intact: **goal met**.
Subjective • Says he has no appetite **Objective** • Braden score 12 • Thin, with bony prominences standing out • Albumin low	**RISK** **Risk for adult pressure injury** as evidenced by risk factors: immobility, moisture, poor nutrition and low Braden score.	No pressure injury develops during his stay.	• Check the skin over bony areas every shift. • Nurse him on a pressure-redistributing mattress. • Unless there is a reason not to, keep the head of the bed at 30 degrees or less. • Refer to the dietitian and encourage protein. • Score the Braden scale again every day.	No pressure injury: **goal met**. Any redness that does not blanch must be reported and documented.