3.17 Obstructive Sleep Apnea

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • His wife says he snores loudly and stops breathing at times • "I feel sleepy all day." **Objective** • BMI 36 kg/m² • Thick neck • SpO_2 falls while he sleeps	**ACTUAL** **Disturbed sleep pattern** related to upper airway obstruction during sleep as evidenced by snoring, apneas and daytime sleepiness.	Within 2 weeks he uses CPAP and says he sleeps better.	• Show him how to use the CPAP mask and look after it. • Encourage him to lose weight. • Tell him to avoid alcohol and sedatives before bed. • Suggest he sleeps on his side.	Uses CPAP every night: **goal met**.
Subjective • "I fell asleep while driving." **Objective** • Sleepy in the day • Dozes off during the interview	**ACTUAL** **Fatigue** related to poor sleep quality as evidenced by daytime sleepiness.	Within 2 weeks of using CPAP, says he feels more alert in the day.	• Until he is treated, he should not drive or use machinery when sleepy. • Encourage a regular sleep routine. • Tell him to avoid sedatives and alcohol in the evening. • Ask about his alertness at follow-up.	Says he is more alert: **goal met**.
Subjective • *No specific complaint* **Objective** • Hypoxemic again and again at night • Obesity	**RISK** **Risk for decreased cardiac tissue perfusion** as evidenced by risk factors: nocturnal hypoxemia and hypertension.	BP stays on target and he uses CPAP every night.	• Check BP at every visit. • Encourage nightly CPAP and taking his antihypertensives. • Encourage weight loss and regular exercise. • Teach him to report chest pain or palpitations.	BP controlled: **goal met**.