3.2 Bronchial Asthma

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "My chest feels tight and I cannot breathe out." **Objective** • Wheezing on breathing out • Expiration is prolonged • RR 28/min, SpO_2 91%	**ACTUAL** **Ineffective breathing pattern** related to bronchoconstriction as evidenced by wheeze, prolonged expiration and tachypnea.	Within 2 hours, RR is 12–20/min and SpO_2 is 94% or more.	• Sit her upright; leaning forward may feel easier. • Give the inhaled short-acting bronchodilator as prescribed and see how she responds. • Give oxygen to keep SpO_2 on target. • Give the corticosteroid as prescribed. • Stay with her and keep calm yourself.	Less wheeze, SpO_2 on target: **goal met**. No better after treatment? Report it at once.
Subjective • "I use the inhaler only when I feel bad." **Objective** • Uses the inhaler the wrong way • Many attacks this year	**ACTUAL** **Ineffective health self-management** related to insufficient knowledge of treatment regimen as evidenced by incorrect inhaler use and frequent exacerbations.	Before discharge she uses the inhaler correctly and can explain how a reliever differs from a preventer.	• Teach inhaler and spacer technique, then watch her do it. • Explain what the reliever does and what the preventer does. • Help her work out her own triggers (dust, smoke, cold air). • Give her a written asthma action plan. • Tell her to rinse her mouth after the steroid inhaler.	Technique correct and explanation accurate: **goal met**.
Subjective • "I wait until the attack is bad before going to hospital." **Objective** • Needs the reliever more than 3 times a week • Has had severe attacks before	**RISK** **Risk for injury** as evidenced by risk factors: delayed treatment of exacerbations and overuse of reliever inhaler.	Can tell you when to follow the action plan and when to go to emergency.	• Explain the warning signs: needing the reliever more often, waking at night, struggling to talk. • Go through the steps of the asthma action plan with her. • If the reliever does not help, she should go to emergency.	Names the warning signs and what to do: **goal met**.