3.6 Pulmonary Tuberculosis

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "Can I stop the tablets when I feel better?" • "I have had a cough for 2 months." **Objective** • Sputum AFB smear positive • Just started anti-TB treatment • Cannot name his medicines	**ACTUAL** **Ineffective health self-management** related to complexity of long-term treatment as evidenced by intention to stop treatment early and inability to name medications.	Before discharge, can give the name, schedule and length of his anti-TB treatment.	• Explain that treatment lasts at least 6 months and must not be stopped early. • Support directly observed therapy (DOT). • Teach the common side effects: orange urine with rifampicin, jaundice, changes in vision, numbness. • Book follow-up sputum tests.	Explains his regimen correctly: **goal met**.
Subjective • "I have no appetite." **Objective** • Lost 5 kg • BMI 17.5 kg/m²	**ACTUAL** **Imbalanced nutrition: less than body requirements** related to anorexia and increased metabolic demand as evidenced by weight loss and low BMI.	Gains weight steadily during treatment.	• Give a high-calorie, high-protein diet. • Give pyridoxine with isoniazid as prescribed. • Weigh once a week.	Weight going up: **goal met**.
Subjective • Lives with a family of 7 **Objective** • Smear-positive TB • Home is crowded	**RISK** **Risk for infection (to others)** as evidenced by risk factors: airborne transmission and close household contacts.	No new cases among contacts; he follows airborne precautions.	• Keep him in airborne isolation until he is no longer infectious. • Teach him to cover his mouth when coughing and to get rid of sputum safely. • Encourage opening windows to air the rooms at home. • Send household contacts for screening.	Follows precautions and contacts have been screened: **goal met**.