4.17 Deep Vein Thrombosis

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "My left calf is swollen and painful." **Objective** • Only one calf swollen • Warm and red • Doppler positive	**ACTUAL** **Ineffective peripheral tissue perfusion** related to venous obstruction as evidenced by swelling, warmth and pain in the affected limb.	Within 3–5 days, swelling and pain are less.	• Give the anticoagulant as prescribed. • Raise the affected leg. • Measure the calf every day. • Never massage the leg. • Encourage walking as allowed.	Swelling down: **goal met**.
Subjective • Gives the pain 6/10 **Objective** • Calf tender • Guards the leg	**ACTUAL** **Acute pain** related to venous inflammation as evidenced by tender calf and reported pain.	Pain 3/10 or less.	• Give analgesia as prescribed; no NSAIDs with anticoagulants unless they are ordered. • Raise the leg. • Put on a compression stocking when ordered.	Pain controlled: **goal met**.
Subjective • *No specific complaint* **Objective** • Clot in a vein	**RISK** **Risk for bleeding** as evidenced by risk factors: anticoagulant therapy.	No bleeding and no sign of pulmonary embolism.	• Check clotting results as ordered. • Look for bleeding from the gums, in the urine and in the stool. • Teach her to report sudden breathlessness or chest pain (could be an embolism). • Watch SpO_2 and RR.	No bleeding, no sign of embolism: **goal met**.