4.3 Angina Pectoris

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I get chest pressure when I walk fast; it goes when I rest." **Objective** • Pain lasts 5 minutes • Eased by nitroglycerin • ST depression on ECG during pain	**ACTUAL** **Acute pain** related to myocardial ischemia as evidenced by exertional chest pressure relieved by rest.	Chest pain relieved within 5–15 minutes of treatment.	• Stop what he is doing and have him sit or lie down. • Give sublingual nitroglycerin as prescribed, checking BP before each dose. • If SpO_2 is below target, give oxygen. • Take a 12-lead ECG while the pain is present. • Pain still there after 3 doses? • Report it (this may be an MI).	Pain relieved: **goal met**.
Subjective • "I do not know how much activity is safe." **Objective** • Newly diagnosed • Cannot explain how to use nitroglycerin	**ACTUAL** **Deficient knowledge** related to new diagnosis as evidenced by questions about activity and medication.	Can explain how to use nitroglycerin and when to get help.	• Teach him to carry nitroglycerin, keep it in a dark container and replace it as directed. • Teach him to call emergency services if the pain is still there after the first dose, following local instructions. • Teach him to build up exercise slowly and not to eat a heavy meal before exertion.	Explains correct use: **goal met**.
Subjective • *No specific complaint* **Objective** • Coronary artery disease • Smokes • Cholesterol high	**RISK** **Risk for decreased cardiac tissue perfusion** as evidenced by risk factors: coronary artery disease, smoking and hyperlipidemia.	Commits to cutting down his risk factors.	• Help him stop smoking. • Teach a low-fat diet. • Encourage him to keep taking his statin and antiplatelet.	Describes his plan to reduce risk: **goal met**.