5.14 Myasthenia Gravis

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "My eyelids droop and I get tired chewing." **Objective** • Eyelids drooping • Chewing and swallowing weak	**ACTUAL** **Impaired swallowing** related to neuromuscular weakness as evidenced by difficulty chewing and swallowing.	Eats safely, without choking.	• Give the anticholinesterase 30–45 minutes before meals, as prescribed. • Offer soft food and let her rest during the meal. • Keep suction available.	Eating safely: **goal met**.
Subjective • *No specific complaint* **Objective** • More tired as the day goes on Eyelids droop more in the evening	**ACTUAL** **Fatigue** related to muscle weakness as evidenced by worsening weakness with activity.	Plans activities for the times when she is strongest.	• Fit activities in soon after her medication. • Encourage rest periods. • Teach ways to save energy.	Activities planned: **goal met**.
Subjective • *No specific complaint* **Objective** • Breathing muscles weak • Cough weak	**RISK** **Risk for aspiration** as evidenced by risk factors: impaired swallowing and weak cough.	No aspiration, and a crisis is recognised early.	• Watch her breathing and swallowing. • Teach the signs of myasthenic and cholinergic crisis.	No aspiration: **goal met**.