6.12 Hernia Repair

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "The groin wound hurts when I move." • Gives the pain 5/10 **Objective** • Incision in the groin • Guards the area when he changes position • HR 96/min	**ACTUAL** **Acute pain** related to surgical incision as evidenced by pain score 5/10 and guarding.	Pain 3/10 or less within 1 hour of analgesia.	• Check pain regularly and after each intervention. • Give analgesia as prescribed. • Show him how to support the wound with a hand or pillow when he coughs or gets up. • If ordered for scrotal swelling, use a scrotal support or ice pack.	Pain 3/10 or less: **goal met**. If not, go over the analgesia with the surgeon.
Subjective • "When can I go back to lifting at work?" **Objective** • Does manual work • Cannot say what he should avoid doing	**ACTUAL** **Deficient knowledge** related to post-operative activity restrictions as evidenced by questions and inability to state limits.	Before discharge, can explain his lifting limits and how to care for the wound.	• He must avoid heavy lifting and straining for as long as the surgeon advises. • Teach him to prevent constipation with fluids and fibre. • Teach wound care and the signs of infection. • Teach him to report a new bulge or severe pain.	States his limits correctly: **goal met**.
Subjective • "I have not passed urine since the operation." **Objective** • Had spinal or general anesthesia • On opioid analgesia • Older man	**RISK** **Risk for urinary retention** as evidenced by risk factors: anesthesia, opioids and groin surgery.	Passes urine within 6–8 hours of surgery.	• Record when he first passes urine and how much. • If he has not voided, feel or scan the bladder. • If allowed, help him stand up to pass urine. • If he still cannot void, catheterise as ordered.	Voids adequately: **goal met**.