6.8 Ulcerative Colitis

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I pass bloody stools 8 times a day." • "I rush to the toilet." **Objective** • Diarrhea with blood • Cramps in the lower abdomen • Hyperactive bowel sounds	**ACTUAL** **Diarrhea** related to inflammation of colonic mucosa as evidenced by frequent bloody stools and urgency.	Within 1 week of treatment, fewer stools with less blood.	• Chart how often, how loose, and whether there is blood. • Give corticosteroids and aminosalicylates as prescribed. • Keep a commode or the toilet close at hand. • Protect the skin around the anus. • She should avoid foods that make things worse.	Fewer stools: **goal met**.
Subjective • "I feel dizzy when I stand." **Objective** • HR 112/min • Hb 9 g/dL • Mouth dry	**ACTUAL** **Deficient fluid volume** related to fluid and blood loss in stool as evidenced by tachycardia, dry mucosa and low Hb.	Within 24 hours, vital signs stable and urine output adequate.	• Give IV fluids and blood as prescribed. • Keep an I&O chart. • Follow potassium and Hb.	Vital signs stable: **goal met**.
Subjective • *No specific complaint* **Objective** • Colitis severe • On high-dose steroids	**RISK** **Risk for injury** as evidenced by risk factors: toxic megacolon and perforation in severe colitis.	Complications are picked up early.	• Measure her abdominal girth every day. • Report sudden distension, fever, a fast heart rate, or fewer stools along with pain. • No antidiarrheals or opioids unless ordered.	No complication: **goal met**.